

# Transcriptional Analysis of Responses of *Desulfovibrio vulgaris* to NaCl Stress Using Whole-Genome Oligonucleotide Microarrays

Zhili He<sup>1</sup>, Qiang He<sup>1</sup>, Liyou Wu<sup>1</sup>, Judy D. Wall<sup>2</sup>, Terry Hazen<sup>3</sup>, Matthew W. Fields<sup>4</sup> and Jizhong Zhou<sup>1</sup>

<sup>1</sup>Oak Ridge National Laboratory, Oak Ridge, TN

<sup>2</sup>University of Missouri, Columbia, MO

<sup>3</sup>Lawrence Berkeley National Laboratory, Berkeley, CA

<sup>4</sup>Miami University, Oxford, OH

The gram-negative sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough is fully sequenced with a genome size of 3.6 Mb and 3584 ORFs. ~~The w~~Whole-genome microarrays of *D. vulgaris* were constructed using 70-mer oligos, containing 97.1% unique probes as well as 2.9% non-specific probes that may have cross-hybridization with other ORFs. The whole genome oligonucleotide microarrays were used to explore mechanisms of *D. vulgaris* to deal with salt stress in its natural habitats.

We first examined ~~its~~ the growth of *D. vulgaris* under different concentrations of NaCl and then investigated the initial reaction to salt shock with 0.5 M NaCl. Growth of *D. vulgaris* cells was not affected significantly under 50 and 100 mM NaCl conditions. Under 250 and 500 mM NaCl conditions, they first grew slowly with about 22-hour and 42-hour lag periods, respectively, and finally they achieved 80% and 50% of the extent [not rate, I assume] of the control growth, respectively. Salt shock with 250 and 500 mM NaCl showed that *D. vulgaris* cells immediately responded to sudden increase in salinity and that their biomass decreased. *D. vulgaris* cells were not able to grow in 1000 mM NaCl medium~~a~~. There were about 370 ORFs up-regulated and around 140 ORFs down-regulated when *D. vulgaris* cells were treated with 0.5 M NaCl for 0.5 hour. For example, ORF02787, ORF02789 and ORF02792 predicted to be genes involved in glycine/betaine/L-proline transportation were up-regulated 5, 19 and 26 fold~~s~~, respectively. As expected, about 50% of those genes significantly changed ~~genes~~ were predicted to be hypothetical proteins. After a 4-hour treatment, approximately 140 ORFs were seen to be up-regulated and more than 700 ORFs<sub>2</sub> to be down-regulated. Patterns of gene expression were distinctly different among different time points.

Further study is in progress on the details of those genes significantly changed in expression ~~genes~~, ~~construction~~ on elucidation of their regulatory networks, and on the progression of monitoring dynamic adaptation of *D. vulgaris* cells to high salinity over time.

Question: Does “whole genome” array include the intergenic sequences? Is this array only ORFs?